**PROCESSED AT :**
**Thyrocare**

MC-5241

Thyrocare
Tests you can trust

*Corporate office* : Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400 703
022 - 3090 0000 / 6712 3400 9870666333 wellness@thyrocare.com www.thyrocare.com

## REPORT

**NAME** : XXXXXXXXXXXXXXXXXXX
**REF. BY** : XXXXXXXXXXXXXXXXXXX
**TEST ASKED** : FOLATE

**HOME COLLECTION :**
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

**PATIENTID** :

| TEST NAME | TECHNOLOGY | VALUE | UNITS |
|---|---|---|---|
| FOLATE | C.L.I.A | 8.3 | ng/mL |

**Reference Range :**
> 5.38 ng/ml

Clinical Significance: Low folate intake, malabsorption as a result of gastrointestinal diseases, pregnancy, and drugs such as phenytoin are causes of folate deficiency.3 Folate deficiency is also associated with chronic alcoholism. Serum folate measurement provides an early index of folate status.

Specifications: Precision: Intra assay (%CV): 7.93, Inter assay (%CV): 7.19, Sensitivity: 0.35 ng/mL.

Kit Validation References: Steinkamp RC. Vitamin B12 and folic acid: clinical and pathophysiological considerations. In: Brewster MA, Naito HK, eds. Nutritional Elements and Clinical Biochemistry. New York: Plenum Publishing Corp.; 1980:169–240

**Method :** COMPETITIVE CHEMI LUMINESCENT IMMUNO ASSAY

**Please correlate with clinical conditions.**

**~~ End of report ~~**

Dummy Report

**Sample Collected on (SCT)** : Sample collection time
**Sample Received on (SRT)** : Sample receiving time at Lab
**Report Released on (RRT)** : Report release time
**Sample Type** : SERUM
**Labcode** :
**Barcode** :

Doctor 1 Sign Doctor 2 Sign