PROCESSED AT :
Thyrocare

Thyrocare
Tests you can trust

Corporate office : Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400 703
022 - 3090 0000 / 6712 3400 9870666333 wellness@thyrocare.com www.thyrocare.com

REPORT

NAME : XXXXXXXXXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXXXXXXXXX
TEST ASKED : HAVM

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

PATIENTID :

TEST NAME	TECHNOLOGY	VALUE	UNITS
ANTI HEPATITIS A VIRUS (ANTI HAV) - IGM	**C.M.I.A**	**4.52**	**S / CO**

Reference Range :-

Negative : < 0.80 (Non Reactive)
Equivocal: 0.80 - 1.20 (Borderline Reactive)
Positive : > 1.20 (Reactive)

Clinical Significance:
During the acute phase of HAV infection, IgM Anti-HAV appears in the patient's serum and is nearly always detectable at the onset of symptoms. In most cases, IgM Anti-HAV response usually peaks within the first month of illness and can persist for up to six months. For diagnostic purpose, results should always be assessed in conjunction with the patients medical history, clinical examination and other findings.

Specifications:
Precision: Intra assay (%CV): 11.33 %, Inter assay (%CV): 13.10%; Specificity of 99.0% and sensitivity of 95.0%

Kit validation references :
Locarnini SA, Ferris AA, Lehmana NL, ET AL. The antibody response following Hepatitis A Infection. Intervirology 1977;8(5): 309-18.

Please correlate with clinical conditions.
Method:- FULLY AUTOMATED CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY

~~ End of report ~~

Dummy Report

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign Doctor 2 Sign