PROCESSED AT :
Thyrocare

Thyrocare
Tests you can trust

Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400703 98706 66333 wellness@thyrocare.com

9 out of 10 Doctors Trust that Thyrocare Reports are Accurate & Reliable

NAME : XXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXX
TEST ASKED : HCVG

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
HEPATITIS C VIRUS - IGG	E.L.I.S.A	6.33	OD Ratio

Bio. Ref. Interval. :-

Negative : < 1.00
Positive : >=1.00

Clinical Significance:
Hepatitis C virus is a small, envelop, positive- sense, single stranded DNA virus. HCV now known to major cause of blood transmitted non- A non- B hepatitis. Antibodies to HCV are detectable about 45 days after exposure. Therefore detection of HCV antibodies in serum or plasma is useful in determination of HCV exposure and in the diagnosis of Hepatitis C.

Specifications:
Sensitivity: 100%, Specificity: 99.81%

Kit validation References:
Wilber J. C. Development and use of laboratory test for hepatitis C infection: a review J. Clin. Immunoassay 1993; 16:20

Please correlate with clinical conditions.
Method:- SOLID PHASE ENZYME IMMUNOASSAY

~~ End of report ~~

Dummy Report

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign Doctor 2 Sign