PROCESSED AT :
Thyrocare

Thyrocare
Tests you can trust

Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400703 98706 66333 wellness@thyrocare.com

*9 out of 10 Doctors Trust that Thyrocare Reports are **Accurate & Reliable***

NAME : XXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXX
TEST ASKED : MEASLES ANTIBODY - IGG

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

PATIENTID : XXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
MEASLES ANTIBODY - IGG	C.L.I.A	< 5	AU/mL

Bio. Ref. Interval. :-

Negative : < 13.5
Equivocal : 13.5 - 16.5
Positive : > 16.5

Clinical Significance:
Measles is acute, highly contagious disease. Measles symptoms appear in two stages. In first stage, the individual may have runny nose, cough and a slight fever. The second stage consist of high fever, red blotchy rash.

Specifications:
Intra assay Precision: 8% (%CV), Inter Assay Precision: 15.8% (%CV), Sensitivity: 94.70%, Specificity: 97.40%

Kit Validation References:
TH.TULCHINSKY et.al measles control in developing and developed countries : the case for a two-dose policy. Bull World health organ ,71(1): 93-103(1993)

Please correlate with clinical conditions.
Method:- SANDWICH CHEMI LUMINESCENT IMMUNO ASSAY

~~ End of report ~~

Dummy Report

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign Doctor 2 Sign