PROCESSED AT :
Thyrocare

Corporate office : **Thyrocare Technologies Limited,** D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400 703
022 - 3090 0000 / 6712 3400 9870666333 wellness@thyrocare.com www.thyrocare.com

REPORT

NAME : XXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXX
TEST ASKED : PROGESTERONE

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
PROGESTERONE	C.M.I.A	5.47	ng/mL

Reference Range :-

Adult males : < 0.10 - 0.20 ng/ml

Normal menstruating females
Follicular phase : < 0.10 - 0.30 ng/ml
Luteal phase : 1.20 - 15.9 ng/ml
Postmenopausal females : < 0.10 - 0.20 ng/ml

Pregnant Women
1st Trimester : 2.80 - 147.3 ng/ml
2nd Trimester : 22.5 - 95.3 ng/ml
3rd Trimester : 27.9 - 242.5 ng/ml

Clinical significance: Clinical evaluation of progesterone confirms ovulation and normal luteal function in nonpregnant women. Inadequate progesterone production by the corpus luteum may indicate luteal phase deficiency (LPD), which is associated with infertility and early miscarriage. For diagnostic purpose, results should always be assessed in conjunction with the patients medical history, clinical examination and other findings.

Specifications: Precision: Intra assay (%CV): 5.5 %, Inter assay (%CV): 6.2%; Sensitivity: < 0.1 ng/ml

Kit Validation Reference: Weigel NL, Rowan BG. Estrogen and progesterone action. In: DeGroot LJ, Jameson JL, et al. eds. Endocrinology. Vol 3. 4th ed. Philadelphia: WB Saunders Co., 2001. 2053-2060

Please correlate with clinical conditions.
Method:- FULLY AUTOMATED CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY

~~ End of report ~~

Dummy Report

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign Doctor 2 Sign