PROCESSED AT :

Thyrocare
Tests you can trust

Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400703 98706 66333 wellness@thyrocare.com

*9 out of 10 Doctors Trust that Thyrocare Reports are **Accurate & Reliable***

NAME : XXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXX
TEST ASKED : UPTR

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

PATIENTID : XXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE
URINE PREGNANCY RAPID TEST	IMMUNOASSAY	NEGATIVE

Clinical Significance :-

This test is based on the qualitative detection of hCG which is secreted by viable placental tissue during pregnancy (upon conceiving). HCG levels appear on the blood and urine after conception and continue rise until 8th week of pregnancy. Positive results should be correlated with physical/clinical conditions with the help of physicians. To cross-verify the negative results (if suspected of pregnancy), the test can be repeated after 48 hours as early pregnancy may not yield detectable hCG levels. HCG levels are also increased in non-pregnant conditions likes trophoblastic and non-trophoblastic neoplasms(tumors).

Kit Validation reference –
Batzer FR."Hormonal evaluations of early pregnancy".Fertil. Steril,1980;34(1):1-13

Please correlate with clinical conditions.
Method:- TWO SITE SANDWICH IMMUNOASSAY

~~ End of report ~~

Dummy Report

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : URINE
Labcode :
Barcode :

Doctor 1 Sign Doctor 2 Sign