**PROCESSED AT :**
**Thyrocare**

Thyrocare
Tests you can trust

Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400703 98706 66333 wellness@thyrocare.com

*9 out of 10 Doctors Trust that Thyrocare Reports are **Accurate & Reliable***

**NAME** : XXXXXXXXXXXXXXXX
**REF. BY** : XXXXXXXXXXXXXXXX
**TEST ASKED** : B2MIC

**SAMPLE COLLECTED AT :**
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

| TEST NAME | TECHNOLOGY | VALUE | UNITS |
|---|---|---|---|
| BETA2 MICROGLOBULIN | C.L.I.A | 1958 | ng/mL |

**Bio. Ref. Interval. :-**

Normal: 609 - 2366 ng/mL

Clinical Significance:

Low levels of beta2-microgloublin can indicate non- progression of HIV. Increased levels may be seen in lymphoproliferative diseases such as multiple myeloma-cell chronic lymphocytic leukemia, Hodgkins disease, non- Hodgkins lymphoma, systemic lupus, rheumatoid arthritis, Crohns disease and certain viral infections. Elevated serum levels have also been observed in some hemodialysis patients and in renal transplant rejection.

Specifications:

Precision: Intra assay (%CV): 4.2, Inter assay (%CV): 10.8, Sensitivity: 4 ng/mL

Kit Validation Reference:

Sims H, Miller L. Beta-2-Microglobulin in laboratory diagnosis. Lab Med 1984;15:803–5.

**Please correlate with clinical conditions.**
**Method:-** SOLID-PHASE TWO-SITE CHEMILUMINESCENT IMMUNOMETRIC ASSAY

**~~ End of report ~~**

Dummy Report

**Sample Collected on (SCT)** : Sample collection time
**Sample Received on (SRT)** : Sample receiving time at Lab
**Report Released on (RRT)** : Report release time
**Sample Type** : SERUM
**Labcode** :
**Barcode** :

Doctor 1 Sign Doctor 2 Sign