Thyrocare

Tests you can trust

Name : XXXXXXXX

Date : XXXXXXXXX

Test Asked : Aarogyam Camp Profile 3

Report Status: Complete Report



9 out of 10 Doctors trust that Thyrocare **reports are accurate & reliable***

98% Reports released within **06 Hours** of sample reaching the lab+

Samples Processed in **NABL Accredited** Labs#

700+ Tests & Profiles

Temperature-Controlled Sample Logistics

Unique Barcode Tracking & Reports with QR Code Verification

Fully Automated Machines Inspected Daily

Abnormal Values Re-Checked Twice

Reports Verified By Expert MD Pathologists Stationed at Every Lab

Accredited by

NABL From 2005#

ISO 9001: 2015 - From 2015

CAP From 2007

Dummy Report

PROCESSED AT :

Thyrocare
Tests you can trust

Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400703 98706 66333 wellness@thyrocare.com

*9 out of 10 Doctors Trust that Thyrocare Reports are **Accurate & Reliable***

NAME : XXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXX
TEST ASKED : AAROGYAM CAMP PROFILE 3

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXX

Report Availability Summary

Note: Please refer to the table below for status of your tests.

17 Ready | **0** Ready with Cancellation | **0** Processing | **0** Cancelled in Lab

TEST DETAILS	REPORT STATUS
AAROGYAM CAMP PROFILE 3	Ready
CHLORIDE	Ready
C-REACTIVE PROTEIN (CRP)	Ready
25-OH VITAMIN D (TOTAL)	Ready
ERYTHROCYTE SEDIMENTATION RATE (ESR)	Ready
HIGH SENSITIVITY C-REACTIVE PROTEIN (HS-CRP)	Ready
Lipoprotein (a) [Lp(a)]	Ready
RHEUMATOID FACTOR (RF)	Ready
SODIUM	Ready
VITAMIN B-12	Ready
HBA PROFILE	Ready
HEMOGRAM - 6 PART (DIFF)	Ready
LIVER FUNCTION TESTS	Ready
IRON DEFICIENCY PROFILE	Ready
KIDPRO	Ready
LIPID PROFILE	Ready
T3-T4-USTSH	Ready
APOLIPROTEIN RATIO	Ready

Dummy Report

PROCESSED AT :

CAP ACCREDITED
COLLEGE of AMERICAN PATHOLOGISTS

Thyrocare
Tests you can trust

Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400703 98706 66333 wellness@thyrocare.com

*9 out of 10 Doctors Trust that Thyrocare Reports are **Accurate & Reliable***

NAME : XXXXXXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXXXXXX
TEST ASKED : AAROGYAM CAMP PROFILE 3

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXX

Summary Report

Tests outside reference range

TEST NAME	OBSERVED VALUE	UNITS	Bio. Ref. Interval.
COMPLETE HEMOGRAM			
HEMATOCRIT(PCV)	33.6	%	40.0-50.0
HEMOGLOBIN	10.4	g/dL	13.0-17.0
LYMPHOCYTE	17.4	%	20-40
MEAN CORP.HEMO.CONC(MCHC)	31	g/dL	31.5-34.5
MEAN CORPUSCULAR HEMOGLOBIN(MCH)	26.8	pq	27.0-32.0
RED CELL DISTRIBUTION WIDTH (RDW-CV)	15.4	%	11.6-14
RED CELL DISTRIBUTION WIDTH - SD(RDW-SD)	48.9	fL	39-46
TOTAL RBC	3.88	X 10^6/µL	4.5-5.5
LIPID			
LDL / HDL RATIO	0.9	Ratio	1.5-3.5
TC/ HDL CHOLESTEROL RATIO	2.2	Ratio	3 - 5
RENAL			
CREATININE - SERUM	0.7	mg/dL	0.72-1.18

Dummy Report

Disclaimer: The above listed is the summary of the parameters with values outside the BRI. For detailed report values, parameter correlation and clinical interpretation, kindly refer to the same in subsequent pages.

PROCESSED AT :

CAP ACCREDITED
COLLEGE of AMERICAN PATHOLOGISTS

Thyrocare
Tests you can trust

Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400703 98706 66333 wellness@thyrocare.com

*9 out of 10 Doctors Trust that Thyrocare Reports are **Accurate & Reliable***

NAME : XXXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXXX
TEST ASKED : ESR,HBA PROFILE,HEMOGRAM

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
HbA1c - (HPLC)			
	H.P.L.C	5.1	%

Bio. Ref. Interval. :

Bio. Ref. Interval.: As per ADA Guidelines	Guidance For Known Diabetics
Below 5.7% : Normal 5.7% - 6.4% : Prediabetic >=6.5% : Diabetic	Below 6.5% : Good Control 6.5% - 7% : Fair Control 7.0% - 8% : Unsatisfactory Control >8% : Poor Control

Method : Fully Automated H.P.L.C method

AVERAGE BLOOD GLUCOSE (ABG)	CALCULATED	100	mg/dL

Bio. Ref. Interval. :

90 - 120 mg/dl : Good Control 121 - 150 mg/dl : Fair Control 151 - 180 mg/dl : Unsatisfactory Control > 180 mg/dl : Poor Control

Method : Derived from HBA1c values

Please correlate with clinical conditions.

Dummy Report

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : EDTA Whole Blood
Labcode :
Barcode :

Doctor 1 Sign Doctor 2 Sign

PROCESSED AT :

CAP ACCREDITED
COLLEGE of AMERICAN PATHOLOGISTS

Thyrocare
Tests you can trust

Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400703 98706 66333 wellness@thyrocare.com

*9 out of 10 Doctors Trust that Thyrocare Reports are **Accurate & Reliable***

NAME : XXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXX
TEST ASKED : ESR,HBA PROFILE,HEMOGRAM

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
ERYTHROCYTE SEDIMENTATION RATE (ESR)	MODIFIED WESTERGREN	< 7.5	mm / hr

Bio. Ref. Interval. :-

Male : 0-15
Female : 0-20

Please correlate with clinical conditions.
Method:- MODIFIED WESTERGREN

Dummy Report

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : EDTA Whole Blood
Labcode :
Barcode :

Doctor 1 Sign Doctor 2 Sign

NAME : XXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXX
TEST ASKED : ESR,HBA PROFILE,HEMOGRAM

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	METHODOLOGY	VALUE	UNITS	Bio. Ref. Interval.
HEMOGLOBIN	**SLS-Hemoglobin Method**	**10.4**	**g/dL**	**13.0-17.0**
Hematocrit (PCV)	**CPH Detection**	**33.6**	**%**	**40.0-50.0**
Total RBC	**HF & EI**	**3.88**	**X 10^6/µL**	**4.5-5.5**
Mean Corpuscular Volume (MCV)	Calculated	86.6	fL	83.0-101.0
Mean Corpuscular Hemoglobin (MCH)	**Calculated**	**26.8**	**pq**	**27.0-32.0**
Mean Corp.Hemo. Conc (MCHC)	**Calculated**	**31**	**g/dL**	**31.5-34.5**
Red Cell Distribution Width - SD (RDW-SD)	**Calculated**	**48.9**	**fL**	**39-46**
Red Cell Distribution Width (RDW - CV)	**Calculated**	**15.4**	**%**	**11.6-14**
RED CELL DISTRIBUTION WIDTH INDEX (RDWI)	Calculated	343.7	-	*Refer Note below
MENTZER INDEX	Calculated	22.3	-	*Refer Note below
TOTAL LEUCOCYTE COUNT (WBC)	HF & FC	7.77	X 10^3 / µL	4.0 - 10.0
DIFFERENTIAL LEUCOCYTE COUNT				
Neutrophils Percentage	Flow Cytometry	75.2	%	40-80
Lymphocytes Percentage	**Flow Cytometry**	**17.4**	**%**	**20-40**
Monocytes Percentage	Flow Cytometry	2.7	%	2-10
Eosinophils Percentage	Flow Cytometry	3.9	%	1-6
Basophils Percentage	Flow Cytometry	0.5	%	0-2
Immature Granulocyte Percentage (IG%)	Flow Cytometry	0.3	%	0-0.5
Nucleated Red Blood Cells %	Flow Cytometry	0.1	%	0.0-5.0
ABSOLUTE LEUCOCYTE COUNT				
Neutrophils - Absolute Count	Calculated	5.84	X 10^3 / µL	2.0-7.0
Lymphocytes - Absolute Count	Calculated	1.35	X 10^3 / µL	1.0-3.0
Monocytes - Absolute Count	Calculated	0.21	X 10^3 / µL	0.2 - 1.0
Basophils - Absolute Count	Calculated	0.04	X 10^3 / µL	0.02 - 0.1
Eosinophils - Absolute Count	Calculated	0.3	X 10^3 / µL	0.02 - 0.5
Immature Granulocytes (IG)	Calculated	0.02	X 10^3 / µL	0-0.3
Nucleated Red Blood Cells	Calculated	0.01	X 10^3 / µL	0.0-0.5
PLATELET COUNT	HF & EI	260	X 10^3 / µL	150-410
Mean Platelet Volume (MPV)	Calculated	10.3	fL	6.5-12
Platelet Distribution Width (PDW)	Calculated	11.2	fL	9.6-15.2
Platelet to Large Cell Ratio (PLCR)	Calculated	26.6	%	19.7-42.4
Plateletcrit (PCT)	Calculated	0.27	%	0.19-0.39

***Note - Mentzer index (MI), RDW-CV and RDWI are hematological indices to differentiate between Iron Deficiency Anemia (IDA) and Beta Thalassemia Trait (BTT). MI >13, RDWI >220 and RDW-CV >14 more likely to be IDA. MI <13, RDWI <220, and RDW-CV <14 more likely to be BTT. Suggested Clinical correlation. BTT to be confirmed with HB electrophoresis if clinically indicated.**

Method : Fully automated bidirectional analyser (6 Part Differential SYSMEX XN-1000)

(Reference : *FC- flowcytometry, *HF- hydrodynamic focussing, *EI- Electric Impedence, *Hb- hemoglobin, *CPH- Cumulative pulse height**)**

Dummy Report

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : EDTA Whole Blood
Labcode :
Barcode :

Doctor 1 Sign Doctor 2 Sign

NAME : XXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXX
TEST ASKED : AAROGYAM CAMP PROFILE 3

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
25-OH VITAMIN D (TOTAL)	C.L.I.A	< 65	ng/mL

Bio. Ref. Interval. :-

DEFICIENCY : <20 ng/ml || INSUFFICIENCY : 20-<30 ng/ml
SUFFICIENCY : 30-100 ng/ml || TOXICITY : >100 ng/ml

Clinical Significance:

Vitamin D is a fat soluble vitamin that has been known to help the body absorb and retain calcium and phosphorous; both are critical for building bone health. Decrease in vitamin D total levels indicate inadequate exposure of sunlight, dietary deficiency, nephrotic syndrome. Increase in vitamin D total levels indicate Vitamin D intoxication.

Specifications: Precision: Intra assay (%CV):5.3%, Inter assay (%CV):11.9% ; Sensitivity:3.2 ng/ml.

Kit Validation Reference: Holick MF. Vitamin D Deficiency. N Engl J Med. 2007;357:266–81.

Please correlate with clinical conditions.
Method:- Fully Automated Chemi Luminescent Immuno Assay

Dummy Report

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign Doctor 2 Sign

NAME : XXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXX
TEST ASKED : AAROGYAM CAMP PROFILE 3

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
RHEUMATOID FACTOR (RF)	IMMUNOTURBIDIMETRY	< 9.01	IU/mL

Bio. Ref. Interval. :
ADULT : <= 18

Clinical Significance:
Rheumatoid factor is an anti IgG autoimmune antibody. There are high concentration of rheumatoid factor in the serum of some disease, especially rheumatoid arthritis patients. It helps to diagnose rheumatism ,systematic lupus erythematosus, chronic hepatitis etc.

Specifications:
Precision %CV :- Intra assay %CV- 1.38% , Inter assay %CV-2.88%, Sensitivity :- 40 IU/mL.

Kit Validation Reference:
Anderson, S.G., Bentzon, M.W., Houba, V. and Krag, P. Bull. Wld. Hlth. Org. 42: 311-318 (1970).

Method : LATEX ENHANCED IMMUNOTURBIDIMETRY

Please correlate with clinical conditions.

Dummy Report

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign Doctor 2 Sign

NAME : XXXXXXXXXXXXXXXXX

REF. BY : XXXXXXXXXXXXXXXXX

TEST ASKED : AAROGYAM CAMP PROFILE 3

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
APOLIPOPROTEIN - A1 (APO-A1)	IMMUNOTURBIDIMETRY	< 119	mg/dL
Bio. Ref. Interval. : Male : 86 - 152 Female : 94 - 162			
Method : FULLY AUTOMATED RATE IMMUNOTURBIDIMETRY – BECKMAN COULTER			
APOLIPOPROTEIN - B (APO-B)	IMMUNOTURBIDIMETRY	< 101	mg/dL
Bio. Ref. Interval. : Male : 56 - 145 Female : 53 - 138			
Method : FULLY AUTOMATED RATE IMMUNOTURBIDIMETRY – BECKMAN COULTER			
APO B / APO A1 RATIO (APO B/A1)	CALCULATED	0.8	Ratio
Bio. Ref. Interval. : Male : 0.40 - 1.26 Female : 0.38 - 1.14			
Method : Derived from serum Apo A1 and Apo B values			

Please correlate with clinical conditions.

Dummy Report

Sample Collected on (SCT) : Sample collection time

Sample Received on (SRT) : Sample receiving time at Lab

Report Released on (RRT) : Report release time

Sample Type : SERUM

Labcode :

Barcode :

Doctor 1 Sign Doctor 2 Sign

NAME : XXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXX
TEST ASKED : AAROGYAM CAMP PROFILE 3

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
HIGH SENSITIVITY C-REACTIVE PROTEIN (HS-CRP)	IMMUNOTURBIDIMETRY	< 1.5	mg/L

Bio. Ref. Interval. :-

< 1.00 - Low Risk
1.00 - 3.00 - Average Risk
>3.00 - 10.00 - High Risk
> 10.00 - Possibly due to Non-Cardiac Inflammation

Disclaimer: Persistent unexplained elevation of HSCRP >10 should be evaluated for non-cardiovascular etiologies such as infection , active arthritis or concurrent illness.

Clinical significance:
High sensitivity C- reactive Protein (HSCRP) can be used as an independent risk marker for the identification of Individuals at risk for future cardiovascular Disease. A coronary artery disease risk assessment should be based on the average of two hs-CRP tests, ideally taken two weeks apart.

Kit Validation Reference:
1.Clinical management of laboratory date in medical practice 2003-3004, 207(2003).
2.Tietz : Textbook of Clinical Chemistry and Molecular diagnostics :Second edition :Chapter 47:Page no.1507- 1508.

Please correlate with clinical conditions.
Method:- FULLY AUTOMATED LATEX AGGLUTINATION – BECKMAN COULTER

Dummy Report

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign Doctor 2 Sign

PROCESSED AT :

Thyrocare
Tests you can trust

Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400703 98706 66333 wellness@thyrocare.com

9 out of 10 Doctors Trust that Thyrocare Reports are Accurate & Reliable

NAME : XXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXX
TEST ASKED : AAROGYAM CAMP PROFILE 3

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
C-REACTIVE PROTEIN (CRP)	IMMUNOTURBIDIMETRY	< 2.5	mg/L

Bio. Ref. Interval. :-

Acute phase determination : < 5 mg/L

Clinical Significance:
It's a protein present in the sera of acutely ill patients that bound cell wall C-polysaccharide of streptococcus pneumoniae and agglutinates the organisms. CRP is one of the strongest acute -phase reactants, with plasma concentrations rising up after myocardial infarction, stress, trauma, infection, inflammation, surgery, or neoplastic proliferation.

Concentrations >5 mg/L suggest the presence of an infection or inflammatory process. Concentrations are generally higher in bacterial than viral infection. The increase in peak is proportional to tissue damage. Determination of CRP is clinically useful to screen activity of inflammatory diseases such as rheumatoid arthritis; SLE;Leukemia;after surgery;to detect rejection in renal allograft recipients;to detect neonatal septicemia and meningitis. However, it is a nonspecific marker and cannot be interpreted without other clinical information.

Reference:
Tietz Textbook of clinical chemistry and molecular diagnosis fifth edition chapter 21 P538-539

Please correlate with clinical conditions.
Method:- FULLY AUTOMATED LATEX AGGLUTINATION – BECKMAN COULTER

Dummy Report

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign Doctor 2 Sign

NAME : XXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXX
TEST ASKED : AAROGYAM CAMP PROFILE 3

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
VITAMIN B-12	C.L.I.A	< 484	pg/mL

Bio. Ref. Interval. :-

Normal : 211 - 911 pg/ml

Clinical significance :
Vitamin B12 or cyanocobalamin, is a complex corrinoid compound found exclusively from animal dietary sources, such as meat, eggs and milk. It is critical in normal DNA synthesis, which in turn affects erythrocyte maturation and in the formation of myelin sheath. Vitamin-B12 is used to find out neurological abnormalities and impaired DNA synthesis associated with macrocytic anemias. For diagnostic purpose, results should always be assessed in conjunction with the patients medical history, clinical examination and other findings.

Specifications: Intra assay (%CV):5.0%, Inter assay (%CV):9.2 %;Sensitivity:45 pg/ml

Kit Validation reference:
Chen IW, Sperling MI, Heminger LA. Vitamin B12. In: Pesce AJ, Kaplan LA, eds. Methods in Clinical Chemistry. St. Louis: CV Mosby; 1987:569–73.

Please correlate with clinical conditions.
Method:- COMPETITIVE CHEMI LUMINESCENT IMMUNO ASSAY

Dummy Report

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign Doctor 2 Sign

NAME : XXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXX
TEST ASKED : AAROGYAM CAMP PROFILE 3

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
Lipoprotein (a) [Lp(a)]	IMMUNOTURBIDIMETRY	< 14.95	mg/dL

Bio. Ref. Interval. :-

Adults : < 30.0 mg/dl

Clinical Significance:
Determination of LPA may be useful to guide management of individuals with a family history of CHD or with existing disease. The levels of LPA in the blood depends on genetic factors; The range of variation in a population is relatively large and hence for diagnostic purpose, results should always be assessed in conjunction with the patient's medical history, clinical examination and other findings.

Specifications:
Precision %CV :- Intra assay %CV- 4.55% , Inter assay %CV-0.86 %

Kit Validation Reference:
Tietz NW,Clinical Guide to Laboratory Tests Philadelphia WB. Saunders 1995 : 442-444

Please correlate with clinical conditions.
Method:- LATEX ENHANCED IMMUNOTURBIDIMETRY

Dummy Report

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign Doctor 2 Sign

NAME : XXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXX
TEST ASKED : AAROGYAM CAMP PROFILE 3

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
IRON **Bio. Ref. Interval. :** Male : 65 - 175 Female : 50 - 170 **Method :** Ferrozine method without deproteinization	PHOTOMETRY	< 120	µg/dL
TOTAL IRON BINDING CAPACITY (TIBC) **Bio. Ref. Interval. :** Male: 225 - 535 µg/dl Female: 215 - 535 µg/dl **Method :** Spectrophotometric Assay	PHOTOMETRY	< 380	µg/dL
% TRANSFERRIN SATURATION **Bio. Ref. Interval. :** 13 - 45 **Method :** Derived from IRON and TIBC values	CALCULATED	32	%
UNSAT.IRON-BINDING CAPACITY(UIBC) **Bio. Ref. Interval. :** 162 - 368 **Method :** SPECTROPHOTOMETRIC ASSAY	PHOTOMETRY	< 265	µg/dL

Please correlate with clinical conditions.

Dummy Report

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign Doctor 2 Sign

NAME : XXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXX
TEST ASKED : AAROGYAM CAMP PROFILE 3

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS	Bio. Ref. Interval.
TOTAL CHOLESTEROL	PHOTOMETRY	123	mg/dL	< 200
HDL CHOLESTEROL - DIRECT	PHOTOMETRY	56	mg/dL	40-60
LDL CHOLESTEROL - DIRECT	PHOTOMETRY	53.5	mg/dL	< 100
TRIGLYCERIDES	PHOTOMETRY	120	mg/dL	< 150
TC/ HDL CHOLESTEROL RATIO	**CALCULATED**	**2.2**	**Ratio**	**3 - 5**
TRIG / HDL RATIO	CALCULATED	2.12	Ratio	< 3.12
LDL / HDL RATIO	**CALCULATED**	**0.9**	**Ratio**	**1.5-3.5**
HDL / LDL RATIO	CALCULATED	1.05	Ratio	> 0.40
NON-HDL CHOLESTEROL	CALCULATED	66.2	mg/dL	< 160
VLDL CHOLESTEROL	CALCULATED	23.94	mg/dL	5 - 40

Please correlate with clinical conditions.

Method :

CHOL - Cholesterol Oxidase, Esterase, Peroxidase
HCHO - Direct Enzymatic Colorimetric
LDL - Direct Measure
TRIG - Enzymatic, End Point
TC/H - Derived from serum Cholesterol and Hdl values
TRI/H - Derived from TRIG and HDL Values
LDL/ - Derived from serum HDL and LDL Values
HD/LD - Derived from HDL and LDL values.
NHDL - Derived from serum Cholesterol and HDL values
VLDL - Derived from serum Triglyceride values

***REFERENCE RANGES AS PER NCEP ATP III GUIDELINES:**

TOTAL CHOLESTEROL	(mg/dl)	HDL	(mg/dl)	LDL	(mg/dl)	TRIGLYCERIDES	(mg/dl)
DESIRABLE	<200	LOW	<40	OPTIMAL	<100	NORMAL	<150
BORDERLINE HIGH	200-239	HIGH	>60	NEAR OPTIMAL	100-129	BORDERLINE HIGH	150-199
HIGH	>240			BORDERLINE HIGH	130-159	HIGH	200-499
				HIGH	160-189	VERY HIGH	>500
				VERY HIGH	>190		

Alert !!! 10-12 hours fasting is mandatory for lipid parameters. If not, values might fluctuate.

Dummy Report

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign Doctor 2 Sign

NAME : XXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXX
TEST ASKED : AAROGYAM CAMP PROFILE 3

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS	Bio. Ref. Interval.
ALKALINE PHOSPHATASE	PHOTOMETRY	< 87	U/L	45-129
BILIRUBIN - TOTAL	PHOTOMETRY	< 0.75	mg/dL	0.3-1.2
BILIRUBIN -DIRECT	PHOTOMETRY	< 0.23	mg/dL	< 0.3
BILIRUBIN (INDIRECT)	CALCULATED	0.52	mg/dL	0-0.9
GAMMA GLUTAMYL TRANSFERASE (GGT)	PHOTOMETRY	< 27.45	U/L	< 55
ASPARTATE AMINOTRANSFERASE (SGOT)	PHOTOMETRY	< 17.5	U/L	< 35
ALANINE TRANSAMINASE (SGPT)	PHOTOMETRY	< 22.5	U/L	< 45
SGOT / SGPT RATIO	CALCULATED	0.78	Ratio	< 2
PROTEIN - TOTAL	PHOTOMETRY	< 6.95	gm/dL	5.7-8.2
ALBUMIN - SERUM	PHOTOMETRY	< 4	gm/dL	3.2-4.8
SERUM GLOBULIN	CALCULATED	< 2.95	gm/dL	2.5-3.4
SERUM ALB/GLOBULIN RATIO	CALCULATED	1.36	Ratio	0.9 - 2

Please correlate with clinical conditions.

Method :

ALKP - Modified IFCC method
BILT - Vanadate Oxidation
BILD - Vanadate Oxidation
BILI - Derived from serum Total and Direct Bilirubin values
GGT - Modified IFCC method
SGOT - IFCC* Without Pyridoxal Phosphate Activation
SGPT - IFCC* Without Pyridoxal Phosphate Activation
OT/PT - Derived from SGOT and SGPT values.
PROT - Biuret Method
SALB - Albumin Bcg[1]method (Colorimetric Assay Endpoint)
SEGB - DERIVED FROM SERUM ALBUMIN AND PROTEIN VALUES
A/GR - Derived from serum Albumin and Protein values

Dummy Report

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode : 2911099817/IT001
Barcode : PEL8SDYA

Doctor 1 Sign Doctor 2 Sign

NAME : XXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXX
TEST ASKED : AAROGYAM CAMP PROFILE 3

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
CHLORIDE	I.S.E	< 102.5	mmol/L

Bio. Ref. Interval. :
ADULTS: 98-107 MMOL/L

Clinical Significance :
An increased level of blood chloride (called hyperchloremia) usually indicates dehydration, but can also occur with other problems that cause high blood sodium, such as Cushing syndrome or kidney disease. Hyperchloremia also occurs when too much base is lost from the body (producing metabolic acidosis) or when a person hyperventilates (causing respiratory alkalosis). A decreased level of blood chloride (called hypochloremia) occurs with any disorder that causes low blood sodium. Hypochloremia also occurs with congestive heart failure, prolonged vomiting or gastric suction, Addison disease, emphysema or other chronic lung diseases (causing respiratory acidosis), and with loss of acid from the body (called metabolic alkalosis).

Method : ION SELECTIVE ELECTRODE - INDIRECT

Please correlate with clinical conditions.

Dummy Report

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign Doctor 2 Sign

NAME : XXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXX
TEST ASKED : AAROGYAM CAMP PROFILE 3

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS	Bio. Ref. Interval.
BLOOD UREA NITROGEN (BUN)	PHOTOMETRY	< 14.01	mg/dL	7.94 - 20.07
SODIUM	I.S.E	< 140.5	mmol/L	136 - 145
CREATININE - SERUM	**PHOTOMETRY**	**0.7**	**mg/dL**	**0.72-1.18**
BUN / Sr.CREATININE RATIO	CALCULATED	20.01	Ratio	9:1-23:1
UREA (CALCULATED)	CALCULATED	29.98	mg/dL	Adult : 17-43
UREA / SR.CREATININE RATIO	CALCULATED	42.83	Ratio	< 52
CALCIUM	PHOTOMETRY	< 9.7	mg/dL	8.8-10.6
URIC ACID	PHOTOMETRY	< 5.8	mg/dL	4.2 - 7.3

Please correlate with clinical conditions.

Method :

BUN - Kinetic UV Assay.
SOD - ION SELECTIVE ELECTRODE - INDIRECT
SCRE - Creatinine Enzymatic Method
B/CR - Derived from serum Bun and Creatinine values
UREAC - Derived from BUN Value.
UR/CR - Derived from UREA and Sr.Creatinine values.
CALC - Arsenazo III Method, End Point.
URIC - Uricase / Peroxidase Method

Dummy Report

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign Doctor 2 Sign

NAME : XXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXX
TEST ASKED : AARAGYAM CAMP PROFILE 3

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS	Bio. Ref. Interval.
TOTAL TRIIODOTHYRONINE (T3)	C.L.I.A	109	ng/dL	60-200
TOTAL THYROXINE (T4)	C.L.I.A	<8.29	µg/dL	4.5-12
TSH - ULTRASENSITIVE	C.M.I.A	<2.92	µIU/mL	0.35-4.94

The Biological Reference Ranges is specific to the age group. Kindly correlate clinically.

Method :

T3 - Competitive Chemi Luminescent Immuno Assay
T4 - Competitive Chemi Luminescent Immuno Assay
USTSH - Fully Automated Chemi Luminescent Microparticle Immunoassay

Disclaimer : Results should always be interpreted using the reference range provided by the laboratory that performed the test. Different laboratories do tests using different technologies, methods and using different reagents which may cause difference. In reference ranges and hence it is recommended to interpret result with assay specific reference ranges provided in the reports. To diagnose and monitor therapy doses, it is recommended to get tested every time at the same Laboratory.

Dummy Report

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign Doctor 2 Sign

PROCESSED AT :

Thyrocare
Tests you can trust

Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400703 98706 66333 wellness@thyrocare.com

9 out of 10 Doctors Trust that Thyrocare Reports are Accurate & Reliable

NAME : XXXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXXX
TEST ASKED : AAROGYAM CAMP PROFILE 3

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
EST. GLOMERULAR FILTRATION RATE (eGFR)	CALCULATED	133	mL/min/1.73 m2

Bio. Ref. Interval. :-

> = 90 : Normal
60 - 89 : Mild Decrease
45 - 59 : Mild to Moderate Decrease
30 - 44 : Moderate to Severe Decrease
15 - 29 : Severe Decrease

Clinical Significance

The normal serum creatinine reference interval does not necessarily reflect a normal GFR for a patient. Because mild and moderate kidney injury is poorly inferred from serum creatinine alone. Thus, it is recommended for clinical laboratories to routinely estimate glomerular filtration rate (eGFR), a "gold standard" measurement for assessment of renal function, and report the value when serum creatinine is measured for patients 18 and older, when appropriate and feasible. It cannot be measured easily in clinical practice, instead, GFR is estimated from equations using serum creatinine, age, race and sex. This provides easy to interpret information for the doctor and patient on the degree of renal impairment since it approximately equates to the percentage of kidney function remaining. Application of CKD-EPI equation together with the other diagnostic tools in renal medicine will further improve the detection and management of patients with CKD.

Reference

Levey AS, Stevens LA, Schmid CH, Zhang YL, Castro AF, 3rd, Feldman HI, et al. A new equation to estimate glomerular filtration rate. Ann Intern Med. 2009;150(9):604-12.

Please correlate with clinical conditions.
Method:- CKD-EPI Creatinine Equation

~~ End of report ~~

Dummy Report

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign Doctor 2 Sign

CONDITIONS OF REPORTING


- The reported results are for information and interpretation of the referring doctor only.
- It is presumed that the tests performed on the specimen belong to the patient; named or identified.
- Results of tests may vary from laboratory to laboratory and also in some parameters from time to time for the same patient.
- Should the results indicate an unexpected abnormality, the same should be reconfirmed.
- Only such medical professionals who understand reporting units, reference ranges and limitations of technologies should interpret results.
- This report is not valid for medico-legal purpose.
- Neither Thyrocare, nor its employees/representatives assume: (a) any liability, responsibility for any loss or damage that may be incurred by any person as a result of presuming the meaning or contents of the report, (b) any claims of any nature whatsoever arising from or relating to the performance of the requested tests as well as any claim for indirect, incidental or consequential damages. The total liability, in any case, of Thyrocare shall not exceed the total amount of invoice for the services provided and paid for.
- Thyrocare Discovery video link :- https://youtu.be/nbdYeRgYyQc

EXPLANATIONS

- Majority of the specimen processed in the laboratory are collected by Pathologists and Hospitals we call them as "Clients".
- **Name** - The name is as declared by the client and recored by the personnel who collected the specimen.
- **Ref.Dr** - The name of the doctor who has recommended testing as declared by the client.
- **Labcode** - This is the accession number in our laboratory and it helps us in archiving and retrieving the data.
- **Barcode** - This is the specimen identity number and it states that the results are for the specimen bearing the barcode (irrespective of the name).
- **SCP** - Specimen Collection Point - This is the location where the blood or specimen was collected as declared by the client.
- **SCT** - Specimen Collection Time - The time when specimen was collected as declared by the client.
- **SRT** - Specimen Receiving Time - This time when the specimen reached our laboratory.
- **RRT** - Report Releasing Time - The time when our pathologist has released the values for Reporting.
- **Reference Range** - Means the range of values in which 95% of the normal population would fall.

SUGGESTIONS

- Values out of reference range requires reconfirmation before starting any medical treatment.
- Retesting is needed if you suspect any quality shortcomings.
- Testing or retesting should be done in accredited laboratories.
- For suggestions, complaints, clinical support or feedback, write to us at **customersupport@thyrocare.com** or call us on **022-3090 0000**





*+T&C Apply, # Upto 95% Samples in NABL Accredited Labs, * As per a survey on doctors' perception of laboratory diagnostics (IJARIIT,2023)*

Dummy Report