Thyrocare

Tests you can trust

Name : xxxxxxxxxxxxxxxx

Date : xxxxxxxxxxxxxxxx

Test Asked : Jaanch Cancer Screening Male Advanced

Report Status: Complete Report



First National Diagnostic Chain to have
100% of its Labs with NABL Accreditation#

98% Reports
released within
06 Hours
of sample reaching the lab+

9 out of 10 Doctors Trust
that Thyrocare
reports are
Accurate & Reliable*

1200+
Tests & Profiles

Temperature-Controlled Sample Logistics

Unique Barcode Tracking

Fully Automated Machines Inspected Daily

Abnormal Values Re-Checked Twice

Reports Verified By Expert MD Pathologists Stationed at Every Lab

Your reports are digitally verifiable

Scan the QR code inside the report to check authenticity of reported values

QR code will remain active for 30 days from report release date

Accredited by

NABL From 2005

ISO 9001: 2015 - From 2015

CAP From 2007

PROCESSED AT :

Thyrocare

Tests you can trust

Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400 703 9870666333 wellness@thyrocare.com

First National Diagnostic Chain to have 100% of its Labs with NABL Accreditation#

NAME : XXXXXXXXXXXXXXXX

REF. BY : XXXXXXXXXXXXXXXX

TEST ASKED : JAANCH CANCER SCREENING MALE ADVANCED

SAMPLE COLLECTED AT :

XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

Report Availability Summary

Note: Please refer to the table below for status of your tests.

5 Ready | **0** Ready with Cancellation | **0** Processing | **0** Cancelled in Lab

TEST DETAILS	REPORT STATUS
JAANCH CANCER SCREENING MALE ADVANCED	Ready
CA 19.9	Ready
CARCINO EMBRYONIC ANTIGEN (CEA)	Ready
PROSTATE PROFILE	Ready
ALPHA FETO PROTEIN	Ready
BETA HCG	Ready

Dummy Report

PROCESSED AT :

CAP ACCREDITED
COLLEGE of AMERICAN PATHOLOGISTS

Thyrocare
Tests you can trust

Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400 703 9870666333 wellness@thyrocare.com

First National Diagnostic Chain to have 100% of its Labs with NABL Accreditation#

NAME : XXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXX
TEST ASKED : JAANCH CANCER SCREENING MALE ADVANCED

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
CA 19.9	C.L.I.A	< 17	U/mL

Bio. Ref. Interval. :-

Adults : < 37.0 U/mL

Clinical Significance:

CA 19-9 is elevated in most patients with advanced Pancreatic Cancer, But it may also be elevated in other cancers, conditions, and diseases such as Colorectal cancer, Lung Cancer, Gall Bladder Cancer, Gallstones, Pancreatitis, Cystic Fibrosis, and Liver Disease. Bile duct obstruction may also cause very high CA 19-9 levels. Very small amounts of CA 19-9 may also be found in healthy patients.

For Diagnostic Purpose, Results should always be assessed in conjunction with the patients medical history, Clinical Examination and other findings.

Specifications:

Intra assay (%CV): 8.71 %, Inter assay(%CV) : 5.00% & Sensitivity: 1.2 U/ml

Kit Validation References:

Steinberg W. The clinical utility of the CA 19.9 Tumor-Associated antigen. AM J Gastroenterol 1990; 85(4): 350

Please correlate with clinical conditions.

Method:- FULLY AUTOMATED TWO STEP SANDWICH IMMUNOASSAY

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign Doctor 2 Sign

Scan QR code to verify authenticity of reported results; active for 30 days from release time.

Dummy Report

PROCESSED AT :

CAP ACCREDITED
COLLEGE of AMERICAN PATHOLOGISTS

Thyrocare
Tests you can trust

Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400 703 9870666333 wellness@thyrocare.com

First National Diagnostic Chain to have 100% of its Labs with NABL Accreditation#

NAME : XXXXXXXXXXXXXXXX

REF. BY : XXXXXXXXXXXXXXXX

TEST ASKED : JAANCH CANCER SCREENING MALE ADVANCED

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
FREE PSA	C.L.I.A	< 0.24	ng/mL

Bio. Ref. Interval. :
< 0.50 ng/ml

Clinical Significance:

Free PSA is the unbound form of Prostate Specific Antigen (PSA). Studies have suggested that the percentage of Free PSA in Total PSA is lower in patients with Prostate Cancer than those with benign Prostate Hyperplasia. The free to Total PSA ratio is now being introduced and studied as an additional tool to help clinician to decide if a patient needs more aggressive evaluation, Such as Prostate Biopsy, to check for prostate cancer. For diagnostic purpose, results should always be assessed in conjunction with the patients medical history, clinical examination and other findings.

Specifications:

Precision: Intra Assay (%CV): 5.3 %, Inter Assay (%CV): 5.2 %;
Sensitivity: 0.01 ng/mL

Kit Validation References

Benson MC, Whang Is, Pantuck A Et Al. Prostate Specific Antigen Density: A means of distinguishing benign prostatic hypertrophy and prostate cancer. J. Urol. 147:815-816, 1992

Method : TWO SITE SANDWICH IMMUNOASSAY

Please correlate with clinical conditions.

Dummy Report

Sample Collected on (SCT) : Sample collection time

Sample Received on (SRT) : Sample receiving time at Lab

Report Released on (RRT) : Report release time

Sample Type : SERUM

Labcode :

Barcode :

Doctor 1 Sign Doctor 2 Sign

PROCESSED AT :

Thyrocare
Tests you can trust

Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400 703 9870666333 wellness@thyrocare.com

First National Diagnostic Chain to have 100% of its Labs with NABL Accreditation#

NAME : XXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXX
TEST ASKED : JAANCH CANCER SCREENING MALE ADVANCED

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
ALPHA FETO PROTEIN	E.C.L.I.A	< 100	IU/mL

Bio. Ref. Interval. :-

Men: 0.5 - 5.5 IU/ml
Non-Pregnant Women: 0.5 - 5.5 IU/ml Pregnancy:
Week Range
14th : 10.41 - 49.40
15th : 13.11 - 57.08
16th : 15.12 - 64.45
17th : 17.72 - 76.11
18th : 19.26 - 91.51
19th : 23.26 - 101.80
20th : 28.05 - 125.85
21st : 33.30 - 92.75

Clinical Significance:
AFP has been used as a cancer marker. AFP testing during pregnancy in combination with Beta HCG and E3, Is recommended as an effective way to determine potential fetal risk of open neural tube defect (NTD).

Specifications: Precision: Intra assay (%CV): 4.1, Inter assay (%CV): 4.2, Sensitivity: 1.5 IU/mL

Refrences : Kaur G, Srivastav J, Sharma S, Huria A, Goel P, Chavan BS. Maternal serum median levels of alpha-foetoprotein, human chorionic gonadotropin & unconjugated estriol in second trimester in pregnant women from north-west India. Indian J Med Res. 2013;138(1):83-8.

Please correlate with clinical conditions.
Method:- SANDWHICH ELECTROCHEMILUMINESCENCE IMMUNOASSAY

Dummy Report

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign Doctor 2 Sign

NAME : XXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXX
TEST ASKED : JAANCH CANCER SCREENING MALE ADVANCED

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
CARCINO EMBRYONIC ANTIGEN (CEA)	C.L.I.A	< 1.25	ng/mL

Bio. Ref. Interval. :-

Non-Smokers : < 2.50 ng/mL
Smokers : < 5.00 ng/mL

Clinical Significance :

CEA is often used to monitor patients with cancers of the gastrointestinal tract (GI). Increased CEA levels can indicate some Non-Cancer related conditions, Such as some forms of inflammation, Cirrhosis, and Peptic Ulcer. Also, Smokers tend to have Higher CEA levels than Non-Smokers. When cancer spreads to other organs, CEA levels rise and may be present in other types of bodily fluids besides blood.

For Diagnostic Purpose, Results should always be assessed in Conjunction with the patients medical history, clinical examination and other findings.

Specifications:

Precision: Intra Assay (%CV): 3.6 %, Inter Assay (%CV): 4.1 %; Sensitivity: 0.5 ng/ml

Kit Validation References:

Statland Be, Winkel P. Neoplasia. In: Kaplan LA, Resc AJ, Editors. Clinical Chemistry, Theory, Analysis and Correlation. 2nd Ed. St. Louis: Cv Mosby, 1989.p 734-5.

Please correlate with clinical conditions.
Method:- FULLY AUTOMATED TWO STEP SANDWICH IMMUNOASSAY

Dummy Report

NAME : XXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXX
TEST ASKED : JAANCH CANCER SCREENING MALE ADVANCED

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
BETA HCG	C.M.I.A	< 100	mIU/mL

Bio. Ref. Interval. :-

Negative : < 10 mIU/ml

Pregnancy:

Week	Range	Week	Range
1st - 2nd	10 - 94	6th - 7th	16380 - 139800
2nd - 3rd	61 - 2922	7th - 11th	12540 - 174600
3rd - 4th	666 - 18900	11th - 16th	3684 - 61800
4th - 5th	1536 - 49380	16th - 21st	2832 - 48060
5th - 6th	13860 - 90600	21st - 39th	1620 - 46860

(Multiply mIU/ml Values By 0.10769 to get ng/ml Values)

Clinical Significance:
Females : The rapid rise in HCG Serum levels after conception makes it an excellent marker for early confirmation and monitoring of pregnancy. HCG levels can be useful in prediction of spontaneous abortions, Aiding in the detection of ectopic pregnancy and multiple gestation. For diagnostic purpose, Results should always be assessed in conjunction with the patients medical history, clinical examination and other findings.

Males and Females : . It may also be found in higher than normal amounts in patients with some types of cancer, including testicular, ovarian, liver, stomach, and lung cancers, and in other disorders. Measuring the amount of beta-hCG in the blood of cancer patients may help to diagnose cancer and find out how well cancer treatment is working. Beta-hCG is a type of tumor marker

Kit Validation References: Braunstein GD, Rasor J, Adler D, Danzer H, Wade Me. Serum Human Chorionic Gonadotropin Levels Throughout Normal Pregnancy. Am J Obstet Gynecol 1976: 126: 678-81.

Please correlate with clinical conditions.
Method:- FULLY AUTOMATED CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY

Dummy Report

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign Doctor 2 Sign

NAME : XXXXXXXXXXXXXXXX
REF. BY : XXXXXXXXXXXXXXXX
TEST ASKED : JAANCH CANCER SCREENING MALE ADVANCED

SAMPLE COLLECTED AT :
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
PROSTATE SPECIFIC ANTIGEN (PSA)	C.L.I.A	< 0.7	ng/mL

Bio. Ref. Interval. :-

Normal : < 4.00 ng/ml
Border line : 4.01 to 10.00 ng/ml

Clinical Significance:

Elevated levels of PSA are associated with prostate cancer, but may also be seen with prostatitis (Inflammation of the prostate) and benign prostatic hyperplasia (BPH). PSA test done along with free PSA provides additional information. Studies have suggested that the percentage of free PSA in total PSA is lower in patients with prostate cancer than those with benign prostate hyperplasia.

Specification:

Precision: Intra assay (%CV): 4.38%, Inter assay (%CV): 4.67%; Sensitivity: 0.01 ng/ml

Kit validation references:

Wang MC, Valenzuala LA, Murphy GP, and Chu TM. Purification of a human prostate-specific antigen. Invest. Urol. 1979; 17: 159

Please correlate with clinical conditions.
Method:- TWO SITE SANDWICH IMMUNOASSAY

Dummy Report

Sample Collected on (SCT) : Sample collection time
Sample Received on (SRT) : Sample receiving time at Lab
Report Released on (RRT) : Report release time
Sample Type : SERUM
Labcode :
Barcode :

Doctor 1 Sign Doctor 2 Sign

PROCESSED AT :

Thyrocare

Tests you can trust

Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400 703 9870666333 wellness@thyrocare.com

First National Diagnostic Chain to have 100% of its Labs with NABL Accreditation#

NAME : XXXXXXXXXXXXXXXX

REF. BY : XXXXXXXXXXXXXXXX

TEST ASKED : JAANCH CANCER SCREENING MALE ADVANCED

SAMPLE COLLECTED AT :

XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
PERCENT FREE PSA	CALCULATED	34.29	%

Bio. Ref. Interval. :-

Normal Range :

greater Than 20.0 %

note :

percent Free Psa Is Used To Determine The Relative Risk Of Prostate Cancer In Individual Men. Probability Of Prostate Cancer For Men With Non Suspicious Dre (digital Rectal Examination) Results And Total Psa Between 4 And 10 Ng/ml, By Patient Age Is :

% Free psa	50-64 Years	65-75 Years
0.00% - 10.00%	56%	55%
10.01% - 15.00%	24%	35%
15.01% - 20.00%	17%	23%
20.01% - 25.00%	10%	20%
> 25.00%	5%	9%

Please correlate with clinical conditions.

Method:- N/A

~~ End of report ~~

Dummy Report

Sample Collected on (SCT) : Sample collection time

Sample Received on (SRT) : Sample receiving time at Lab

Report Released on (RRT) : Report release time

Sample Type : SERUM

Labcode :

Barcode :

Doctor 1 Sign Doctor 2 Sign

Scan QR code to verify authenticity of reported results; active for 30days from release time.

CONDITIONS OF REPORTING

- v The reported results are for information and interpretation of the referring doctor only.
- v It is presumed that the tests performed on the specimen belong to the patient; named or identified.
- v Results of tests may vary from laboratory to laboratory and also in some parameters from time to time for the same patient.
- v Should the results indicate an unexpected abnormality, the same should be reconfirmed.
- v Only such medical professionals who understand reporting units, reference ranges and limitations of technologies should interpret results.
- v This report is not valid for medico-legal purpose.
- v Neither Thyrocare, nor its employees/representatives assume: (a) any liability, responsibility for any loss or damage that may be incurred by any person as a result of presuming the meaning or contents of the report, (b) any claims of any nature whatsoever arising from or relating to the performance of the requested tests as well as any claim for indirect, incidental or consequential damages. The total liability, in any case, of Thyrocare shall not exceed the total amount of invoice for the services provided and paid for.
- v Thyrocare Discovery video link :- https://youtu.be/nbdYeRgYyQc

EXPLANATIONS

- v Majority of the specimen processed in the laboratory are collected by Pathologists and Hospitals we call them as "Clients".
- v **Name** - The name is as declared by the client and recored by the personnel who collected the specimen.
- v **Ref.Dr** - The name of the doctor who has recommended testing as declared by the client.
- v **Labcode** - This is the accession number in our laboratory and it helps us in archiving and retrieving the data.
- v **Barcode** - This is the specimen identity number and it states that the results are for the specimen bearing the barcode (irrespective of the name).
- v **SCP** - Specimen Collection Point - This is the location where the blood or specimen was collected as declared by the client.
- v **SCT** - Specimen Collection Time - The time when specimen was collected as declared by the client.
- v **SRT** - Specimen Receiving Time - This time when the specimen reached our laboratory.
- v **RRT** - Report Releasing Time - The time when our pathologist has released the values for Reporting.
- v **Reference Range** - Means the range of values in which 95% of the normal population would fall.

SUGGESTIONS

- v Values out of reference range requires reconfirmation before starting any medical treatment.
- v Retesting is needed if you suspect any quality shortcomings.
- v Testing or retesting should be done in accredited laboratories.
- v For suggestions, complaints, clinical support or feedback, write to us at **customersupport@thyrocare.com** or call us on **022-3090 0000**

Preventive Healthcare is now at your fingertips!

Explore & Select Test / Profile | Book Through App | Booking Confirmation | Track your Technician | Blood Collection | Sample Testing | Download Report & Receipt

Download Thyroapp Now

Launching

Jaanch

For a closer look at your health with



Doctor-Curated Specialised Packages All Under One Roof

Heart | Fever | Cancer | Women's - Reproductive Health

Thyroid | Diabetes | STDs | Skin Care | Hair Fall

Thyrocare
Tests you can trust

*+T&C Apply, #As on 5th December 2024, *As per a survey on doctors' perception of laboratory diagnostics (IJARIIT,2023)*