Thyrocare

Tests you can trust

Name : XXXXXXXXXXXXXXXX

Date : XXXXXXXXX

Test Asked : Jaanch Diabetes Advanced

Report Status : Complete Report

Dummy Report







Thyrocare
Tests you can trust

Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400 703. 9870666333 wellness@thyrocare.com

First National Diagnostic Chain to have 100% of its Labs with NABL Accreditation#

Patient Name : XXXXXXXXXXXX
Referred By : XXXXXXXXXXXX
Address : XXXXXXXXXXXX

Tests Done : JAANCH DIABETES ADVANCED

Report Availability Summary

Note: Please refer to the table below for status of your tests.

9 Ready | **0** Ready with Cancellation | **0** Processing | **0** Cancelled in Lab

TEST DETAILS	REPORT STATUS
JAANCH DIABETES ADVANCED	Ready
FASTING BLOOD SUGAR(GLUCOSE)	Ready
CHLORIDE	Ready
SODIUM	Ready
URINARY MICROALBUMIN	Ready
HBA PROFILE	Ready
HEMOGRAM - 6 PART (DIFF)	Ready
ROUTINE URINE ANALYSIS	Ready
KIDPRO	Ready
LIPID PROFILE	Ready

Dummy Report

Patient Name	: XXXXXXXXXXXX	Tests Done	: JAANCH DIABETES ADVANCED
Referred By	: XXXXXXXXXXXX		
Address	: XXXXXXXXXXXX		

Tests Outside Reference Range

Note: Please refer to the table below for tests outside reference range.

Test Name	Observed Value	Units	Bio. Ref. Interval.
COMPLETE HEMOGRAM			
HEMATOCRIT(PCV)	**33.6**	%	40.0-50.0
HEMOGLOBIN	**10.4**	g/dL	13.0-17.0
LYMPHOCYTE	**17.4**	%	20-40
MEAN CORP.HEMO.CONC(MCHC)	**31**	g/dL	31.5-34.5
MEAN CORPUSCULAR HEMOGLOBIN(MCH)	**26.8**	pq	27.0-32.0
RED CELL DISTRIBUTION WIDTH (RDW-CV)	**15.4**	%	11.6-14
RED CELL DISTRIBUTION WIDTH - SD(RDW-SD)	**48.9**	fL	39-46
TOTAL RBC	**3.88**	X 10^6/µL	4.5-5.5
LIPID			
LDL / HDL RATIO	**0.9**	Ratio	1.5-3.5
TC/ HDL CHOLESTEROL RATIO	**2.2**	Ratio	3 - 5
RENAL			
CREATININE - SERUM	**0.7**	mg/dL	0.72-1.18
URINOGRAM			
SPECIFIC GRAVITY	**< 1.003**	-	1.003-1.030

Dummy Report

Disclaimer: The above listed is the summary of the parameters with values outside the BRI. For detailed report values, parameter correlation and clinical interpretation, kindly refer to the same in subsequent pages.

Thyrocare Technologies Limited, D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400 703. 9870666333 wellness@thyrocare.com

First National Diagnostic Chain to have 100% of its Labs with NABL Accreditation#

Patient Name	: XXXXXXXXXXXX	Sample Collected on (SCT)	: XXXXXXXXXXXX
Referred By	: XXXXXXXXXXXX	Sample Received on (SRT)	: XXXXXXXXXXXX
Address	: XXXXXXXXXXXX	Report Released on (RRT)	: XXXXXXXXXXXX
		Sample Type | Barcode	: XXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
DIABETES SCREEN (URINE)			
URINARY MICROALBUMIN	PHOTOMETRY	< 12.46	µg/mL
Bio. Ref. Interval. : Adults: Less than 25 µg/ml			
Method : Fully Automated Immuno Turbidometry			
CREATININE - URINE	PHOTOMETRY	< 149	mg/dL
Bio. Ref. Interval. : Male: 39 - 259 mg/dl Female: 28 - 217 mg/dl			
Method : Creatinine Jaffe Method, Rate-Blanked and Compensated			
URI. ALBUMIN/CREATININE RATIO (UA/C)	CALCULATED	8.4	µg/mg of Creatinine
Bio. Ref. Interval. : Adults : Less than 30 µg/mg of Creatinine			
Method : Derived from Albumin and Creatinine values			

Please correlate with clinical conditions.

Dummy Report

Tests Done : JAANCH DIABETES ADVANCED

Doctor 1 Sign **Doctor 2 Sign**

Report Remarks : Labcode:2005124167/IT011

Patient Name	: XXXXXXXXXXXX	Sample Collected on (SCT)	: XXXXXXXXXXXX
Referred By	: XXXXXXXXXXXX	Sample Received on (SRT)	: XXXXXXXXXXXX
Address	: XXXXXXXXXXXX	Report Released on (RRT)	: XXXXXXXXXXXX
		Sample Type | Barcode	: XXXXXXXXXXXX

TEST NAME	METHODOLOGY	VALUE	UNITS	Bio. Ref. Interval.
Complete Urinogram				
Physical Examination				
VOLUME	Visual Determination	1	mL	-
COLOUR	Visual Determination	PALE YELLOW	-	Pale Yellow
APPEARANCE	Visual Determination	CLEAR	-	Clear
SPECIFIC GRAVITY	**pKa change**	**< 1.003**	**-**	**1.003-1.030**
PH	pH indicator	5	-	5-8
Chemical Examination				
URINARY PROTEIN	PEI	ABSENT	mg/dL	Absent
URINARY GLUCOSE	GOD-POD	ABSENT	mg/dL	Absent
URINE KETONE	Nitroprusside	ABSENT	mg/dL	Absent
URINARY BILIRUBIN	Diazo coupling	ABSENT	mg/dL	Absent
UROBILINOGEN	Diazo coupling	Normal	mg/dL	<=0.2
URINE BLOOD	Peroxidase reaction	ABSENT	-	Absent
NITRITE	Diazo coupling	ABSENT	-	Absent
LEUCOCYTE ESTERASE	Esterase reaction	ABSENT	-	Absent
Microscopic Examination				
URINARY LEUCOCYTES (PUS CELLS)	Microscopy	2.5	cells/HPF	0-5

(Reference : *PEI - Protein error of indicator, *GOD-POD - Glucose oxidase-peroxidase**)**

Dummy Report

Tests Done : ROUTINE URINE ANALYSIS,URINARY MICROALBUMIN

Doctor 1 Sign **Doctor 2 Sign**

Report Remarks : Labcode:2005124167/IT011

Patient Name	: XXXXXXXXXXXX	Sample Collected on (SCT)	: XXXXXXXXXXXX
Referred By	: XXXXXXXXXXXX	Sample Received on (SRT)	: XXXXXXXXXXXX
Address	: XXXXXXXXXXXX	Report Released on (RRT)	: XXXXXXXXXXXX
		Sample Type | Barcode	: XXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS	Bio. Ref. Interval.
TOTAL CHOLESTEROL	PHOTOMETRY	123	mg/dL	< 200
HDL CHOLESTEROL - DIRECT	PHOTOMETRY	56	mg/dL	40-60
LDL CHOLESTEROL - DIRECT	PHOTOMETRY	53.5	mg/dL	< 100
TRIGLYCERIDES	PHOTOMETRY	120	mg/dL	< 150
TC/ HDL CHOLESTEROL RATIO	**CALCULATED**	**2.2**	**Ratio**	**3 - 5**
TRIG / HDL RATIO	CALCULATED	2.12	Ratio	< 3.12
LDL / HDL RATIO	**CALCULATED**	**0.9**	**Ratio**	**1.5-3.5**
HDL / LDL RATIO	CALCULATED	1.05	Ratio	> 0.40
NON-HDL CHOLESTEROL	CALCULATED	66.2	mg/dL	< 160
VLDL CHOLESTEROL	CALCULATED	23.94	mg/dL	5 - 40

Please correlate with clinical conditions.

Method :

CHOL - Cholesterol Oxidase, Esterase, Peroxidase
HCHO - Direct Enzymatic Colorimetric
LDL - Direct Measure
TRIG - Enzymatic, End Point
TC/H - Derived from serum Cholesterol and Hdl values
TRI/H - Derived from TRIG and HDL Values
LDL/ - Derived from serum HDL and LDL Values
HD/LD - Derived from HDL and LDL values.
NHDL - Derived from serum Cholesterol and HDL values
VLDL - Derived from serum Triglyceride values

***REFERENCE RANGES AS PER NCEP ATP III GUIDELINES:**

TOTAL CHOLESTEROL	(mg/dl)	HDL	(mg/dl)	LDL	(mg/dl)	TRIGLYCERIDES	(mg/dl)
DESIRABLE	<200	LOW	<40	OPTIMAL	<100	NORMAL	<150
BORDERLINE HIGH	200-239	HIGH	>60	NEAR OPTIMAL	100-129	BORDERLINE HIGH	150-199
HIGH	>240			BORDERLINE HIGH	130-159	HIGH	200-499
				HIGH	160-189	VERY HIGH	>500
				VERY HIGH	>190		

Alert !!! 10-12 hours fasting is mandatory for lipid parameters. If not, values might fluctuate.

Dummy Report

Tests Done : JAANCH DIABETES ADVANCED

Report Remarks : Labcode:2005124168/IT011

Doctor 1 Sign

Doctor 2 Sign

Patient Name	: XXXXXXXXXXXX	**Sample Collected on (SCT)**	: XXXXXXXXXXXX
Referred By	: XXXXXXXXXXXX	**Sample Received on (SRT)**	: XXXXXXXXXXXX
Address	: XXXXXXXXXXXX	**Report Released on (RRT)**	: XXXXXXXXXXXX
		Sample Type | Barcode	: XXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
CHLORIDE	I.S.E - INDIRECT	101	mmol/L

Bio. Ref. Interval. :
ADULTS: 98-107 MMOL/L

Clinical Significance :
An increased level of blood chloride (called hyperchloremia) usually indicates dehydration, but can also occur with other problems that cause high blood sodium, such as Cushing syndrome or kidney disease. Hyperchloremia also occurs when too much base is lost from the body (producing metabolic acidosis) or when a person hyperventilates (causing respiratory alkalosis). A decreased level of blood chloride (called hypochloremia) occurs with any disorder that causes low blood sodium. Hypochloremia also occurs with congestive heart failure, prolonged vomiting or gastric suction, Addison disease, emphysema or other chronic lung diseases (causing respiratory acidosis), and with loss of acid from the body (called metabolic alkalosis).

Method : ION SELECTIVE ELECTRODE - INDIRECT

SODIUM	I.S.E - INDIRECT	137	mmol/L

Bio. Ref. Interval. :
Adults: 136-145 mmol/l

Method : ION SELECTIVE ELECTRODE - INDIRECT

Please correlate with clinical conditions.

Dummy Report

Tests Done : JAANCH DIABETES ADVANCED

Doctor 1 Sign **Doctor 2 Sign**

Report Remarks : Labcode:2005124168/IT011

Patient Name	: XXXXXXXXXXXX	Sample Collected on (SCT)	: XXXXXXXXXXXX
Referred By	: XXXXXXXXXXXX	Sample Received on (SRT)	: XXXXXXXXXXXX
Address	: XXXXXXXXXXXX	Report Released on (RRT)	: XXXXXXXXXXXX
		Sample Type | Barcode	: XXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS	Bio. Ref. Interval.
BLOOD UREA NITROGEN (BUN)	PHOTOMETRY	12.6	mg/dL	7.94 - 20.07
CREATININE - SERUM	**PHOTOMETRY**	**0.7**	**mg/dL**	**0.72-1.18**
BUN / Sr.CREATININE RATIO	CALCULATED	18	Ratio	9:1-23:1
UREA (CALCULATED)	CALCULATED	26.96	mg/dL	Adult : 17-43
UREA / SR.CREATININE RATIO	CALCULATED	38.52	Ratio	< 52
CALCIUM	PHOTOMETRY	9.62	mg/dL	8.8-10.6
URIC ACID	PHOTOMETRY	6.8	mg/dL	4.2 - 7.3

Please correlate with clinical conditions.

Method :

BUN - Kinetic UV Assay.
SCRE - Creatinine Enzymatic Method
B/CR - Derived from serum Bun and Creatinine values
UREAC - Derived from BUN Value.
UR/CR - Derived from UREA and Sr.Creatinine values.
CALC - Arsenazo III Method, End Point.
URIC - Uricase / Peroxidase Method

Dummy Report

Tests Done : JAANCH DIABETES ADVANCED

Report Remarks : Labcode:2005124168/IT011

Doctor 1 Sign

Doctor 2 Sign

Patient Name	: XXXXXXXXXXXX	Sample Collected on (SCT)	: XXXXXXXXXXXX
Referred By	: XXXXXXXXXXXX	Sample Received on (SRT)	: XXXXXXXXXXXX
Address	: XXXXXXXXXXXX	Report Released on (RRT)	: XXXXXXXXXXXX
		Sample Type | Barcode	: XXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
EST. GLOMERULAR FILTRATION RATE (eGFR)	CALCULATED	134	mL/min/1.73 m2

Bio. Ref. Interval. :-

> = 90 : Normal
60 - 89 : Mild Decrease
45 - 59 : Mild to Moderate Decrease
30 - 44 : Moderate to Severe Decrease
15 - 29 : Severe Decrease
<15 : Kidney Failure

Clinical Significance

The normal serum creatinine reference interval does not necessarily reflect a normal GFR for a patient. Because mild and moderate kidney injury is poorly inferred from serum creatinine alone. Thus, it is recommended for clinical laboratories to routinely estimate glomerular filtration rate (eGFR), a "gold standard" measurement for assessment of renal function, and report the value when serum creatinine is measured for patients 18 and older, when appropriate and feasible. It cannot be measured easily in clinical practice, instead, GFR is estimated from equations using serum creatinine, age, race and sex. This provides easy to interpret information for the doctor and patient on the degree of renal impairment since it approximately equates to the percentage of kidney function remaining. Application of CKD-EPI equation together with the other diagnostic tools in renal medicine will further improve the detection and management of patients with CKD.

Reference

Levey AS, Stevens LA, Schmid CH, Zhang YL, Castro AF, 3rd, Feldman HI, et al. A new equation to estimate glomerular filtration rate. Ann Intern Med. 2009;150(9):604-12.

Please correlate with clinical conditions.

Method:- 2021 CKD EPI Creatinine Equation

Dummy Report

Tests Done : JAANCH DIABETES ADVANCED

Doctor 1 Sign **Doctor 2 Sign**

Report Remarks : Labcode:2005124168/IT011

Patient Name	: XXXXXXXXXXXX	Sample Collected on (SCT)	: XXXXXXXXXXXX
Referred By	: XXXXXXXXXXXX	Sample Received on (SRT)	: XXXXXXXXXXXX
Address	: XXXXXXXXXXXX	Report Released on (RRT)	: XXXXXXXXXXXX
		Sample Type | Barcode	: XXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
HbA1c	H.P.L.C	5.1	%

Bio. Ref. Interval. :

As per ADA Guidelines	Guidance For Known Diabetics
Below 5.7% : Normal	Below 6.5% : Good Control
5.7% - 6.4% : Prediabetic	6.5% - 7% : Fair Control
>=6.5% : Diabetic	7.0% - 8% : Unsatisfactory Control
	>8% : Poor Control

Method : Fully Automated H.P.L.C method

AVERAGE BLOOD GLUCOSE (ABG)	CALCULATED	100	mg/dL

Bio. Ref. Interval. :

90 - 120 mg/dl : Good Control
121 - 150 mg/dl : Fair Control
151 - 180 mg/dl : Unsatisfactory Control
> 180 mg/dl : Poor Control

Method : Derived from HBA1c values

Please correlate with clinical conditions.

Dummy Report

Tests Done : HBA PROFILE,HEMOGRAM

Doctor 1 Sign **Doctor 2 Sign**

Report Remarks : Labcode:2005124169/IT011

Patient Name	: XXXXXXXXXXXX	Sample Collected on (SCT)	: XXXXXXXXXXXX
Referred By	: XXXXXXXXXXXX	Sample Received on (SRT)	: XXXXXXXXXXXX
Address	: XXXXXXXXXXXX	Report Released on (RRT)	: XXXXXXXXXXXX
		Sample Type | Barcode	: XXXXXXXXXXXX

TEST NAME	METHODOLOGY	VALUE	UNITS	Bio. Ref. Interval.
HEMOGLOBIN	**SLS-Hemoglobin Method**	**10.4**	**g/dL**	**13.0-17.0**
Hematocrit (PCV)	**CPH Detection**	**33.6**	**%**	**40.0-50.0**
Total RBC	**HF & EI**	**3.88**	**X 10^6/µL**	**4.5-5.5**
Mean Corpuscular Volume (MCV)	Calculated	86.6	fL	83.0-101.0
Mean Corpuscular Hemoglobin (MCH)	**Calculated**	**26.8**	**pq**	**27.0-32.0**
Mean Corp.Hemo. Conc (MCHC)	**Calculated**	**31**	**g/dL**	**31.5-34.5**
Red Cell Distribution Width - SD (RDW-SD)	**Calculated**	**48.9**	**fL**	**39-46**
Red Cell Distribution Width (RDW - CV)	**Calculated**	**15.4**	**%**	**11.6-14**
RED CELL DISTRIBUTION WIDTH INDEX (RDWI)	Calculated	343.7	-	*Refer Note below
MENTZER INDEX	Calculated	22.3	-	*Refer Note below
TOTAL LEUCOCYTE COUNT (WBC)	HF & FC	7.77	X 10^3 / µL	4.0 - 10.0
DIFFERENTIAL LEUCOCYTE COUNT				
Neutrophils Percentage	Flow Cytometry	75.2	%	40-80
Lymphocytes Percentage	**Flow Cytometry**	**17.4**	**%**	**20-40**
Monocytes Percentage	Flow Cytometry	2.7	%	2-10
Eosinophils Percentage	Flow Cytometry	3.9	%	1-6
Basophils Percentage	Flow Cytometry	0.5	%	0-2
Immature Granulocyte Percentage (IG%)	Flow Cytometry	0.3	%	0-0.5
Nucleated Red Blood Cells %	Flow Cytometry	0.1	%	0.0-5.0
ABSOLUTE LEUCOCYTE COUNT				
Neutrophils - Absolute Count	Calculated	5.84	X 10^3 / µL	2.0-7.0
Lymphocytes - Absolute Count	Calculated	1.35	X 10^3 / µL	1.0-3.0
Monocytes - Absolute Count	Calculated	0.21	X 10^3 / µL	0.2 - 1.0
Basophils - Absolute Count	Calculated	0.04	X 10^3 / µL	0.02 - 0.1
Eosinophils - Absolute Count	Calculated	0.3	X 10^3 / µL	0.02 - 0.5
Immature Granulocytes (IG)	Calculated	0.02	X 10^3 / µL	0-0.3
Nucleated Red Blood Cells	Calculated	0.01	X 10^3 / µL	0.0-0.5
PLATELET COUNT	HF & EI	260	X 10^3 / µL	150-410
Mean Platelet Volume (MPV)	Calculated	10.3	fL	6.5-12
Platelet Distribution Width (PDW)	Calculated	11.2	fL	9.6-15.2
Platelet to Large Cell Ratio (PLCR)	Calculated	26.6	%	19.7-42.4
Plateletcrit (PCT)	Calculated	0.27	%	0.19-0.39

Remarks : Alert!!! RBCs:Mild anisopoikilocytosis. Predominantly normocytic normochromic with ovalocytes. Platelets:Appear adequate in smear.

***Note - Mentzer index (MI), RDW-CV and RDWI are hematological indices to differentiate between Iron Deficiency Anemia(IDA) and Beta Thalassemia Trait (BTT). MI >13, RDWI >220 and RDW-CV >14 more likely to be IDA. MI <13, RDWI <220, and RDW-CV <14 more likely to be BTT. Suggested Clinical correlation. BTT to be confirmed with HB electrophoresis if clinically indicated.**

Method : Fully automated bidirectional analyser

(Reference : *FC- flowcytometry, *HF- hydrodynamic focussing, *EI- Electric Impedence, *Hb- hemoglobin, *CPH- Cumulative pulse height)

Tests Done : HBA PROFILE,HEMOGRAM

Doctor 1 Sign **Doctor 2 Sign**

Report Remarks : Labcode:2005124169/IT011

Dummy Report

Patient Name	: XXXXXXXXXXXX	Sample Collected on (SCT)	: XXXXXXXXXXXX
Referred By	: XXXXXXXXXXXX	Sample Received on (SRT)	: XXXXXXXXXXXX
Address	: XXXXXXXXXXXX	Report Released on (RRT)	: XXXXXXXXXXXX
		Sample Type | Barcode	: XXXXXXXXXXXX

TEST NAME	TECHNOLOGY	VALUE	UNITS
FASTING BLOOD SUGAR(GLUCOSE)	PHOTOMETRY	90.1	mg/dL

Bio. Ref. Interval. :-

As per ADA Guideline: Fasting Plasma Glucose (FPG)	
Normal	70 to 100 mg/dl
Prediabetes	100 mg/dl to 125 mg/dl
Diabetes	126 mg/dl or higher

Note :
The assay could be affected mildly and may result in anomalous values if serum samples have heterophilic antibodies, hemolyzed , icteric or lipemic. The concentration of Glucose in a given specimen may vary due to differences in assay methods, calibration and reagent specificity. For diagnostic purposes results should always be assessed in conjunction with patients medical history, clinical findings and other findings.

Please correlate with clinical conditions.

Method:- GOD-POD METHOD

~~ End of report ~~

Dummy Report

Tests Done : BLOOD SUGAR (F)

Doctor 1 Sign **Doctor 2 Sign**

Report Remarks : Labcode:2005124170/IT011

CONDITIONS OF REPORTING

- The reported results are for information and interpretation of the referring doctor only.
- It is presumed that the tests performed on the specimen belong to the patient; named or identified.
- Results of tests may vary from laboratory to laboratory and also in some parameters from time to time for the same patient.
- Should the results indicate an unexpected abnormality, the same should be reconfirmed.
- Only such medical professionals who understand reporting units, reference ranges and limitations of technologies should interpret results.
- This report is not valid for medico-legal purpose.
- Neither Thyrocare, nor its employees/representatives assume: (a) any liability, responsibility for any loss or damage that may be incurred by any person as a result of presuming the meaning or contents of the report, (b) any claims of any nature whatsoever arising from or relating to the performance of the requested tests as well as any claim for indirect, incidental or consequential damages. The total liability, in any case, of Thyrocare shall not exceed the total amount of invoice for the services provided and paid for.
- Thyrocare Discovery video link :- https://youtu.be/nbdYeRgYyQc

EXPLANATIONS

- Majority of the specimen processed in the laboratory are collected by Pathologists and Hospitals we call them as "Clients".
- **Name** - The name is as declared by the client and recored by the personnel who collected the specimen.
- **Ref.Dr** - The name of the doctor who has recommended testing as declared by the client.
- **Labcode** - This is the accession number in our laboratory and it helps us in archiving and retrieving the data.
- **Barcode** - This is the specimen identity number and it states that the results are for the specimen bearing the barcode (irrespective of the name).
- **SCP** - Specimen Collection Point - This is the location where the blood or specimen was collected as declared by the client.
- **SCT** - Specimen Collection Time - The time when specimen was collected as declared by the client.
- **SRT** - Specimen Receiving Time - This time when the specimen reached our laboratory.
- **RRT** - Report Releasing Time - The time when our pathologist has released the values for Reporting.
- **Reference Range** - Means the range of values in which 95% of the normal population would fall.

SUGGESTIONS

- Values out of reference range requires reconfirmation before starting any medical treatment.
- Retesting is needed if you suspect any quality shortcomings.
- Testing or retesting should be done in accredited laboratories.
- For suggestions, complaints, clinical support or feedback, write to us at **customersupport@thyrocare.com** or call us on **022-3090 0000**





+ *T&C Apply, #As on 5th December 2024 (Applicable for all company owned labs except Bhagalpur & Vijayawada),*

⋆ *As per survey on doctors' perception of laboratory diagnostics (IJARIIT, 2023), -Mumbai Reference Lab is CAP Accredited*

Dummy Report